

## Half-Life

**Unit:** Atomic and Nuclear Physics

**MA Curriculum Frameworks (2016):** N/A

**AP Physics 2 Learning Objectives:** 7.C.3.1

**Mastery Objective(s):** (Students will be able to...)

- Calculate the amount of material remaining after an amount of time.
- Calculate the elapsed time based on the amount of material remaining.

**Success Criteria:**

- Variables are correctly identified and substituted correctly into the correct equation.
- Algebra is correct and rounding to appropriate number of significant figures is reasonable.

**Language Objectives:**

- Explain why the *mass* of material that decays keeps decreasing.

**Tier 2 Vocabulary:** life, decay

**Labs, Activities & Demonstrations:**

- half-life of dice or M & M candies

**Notes:**

The atoms of radioactive elements are unstable, and they spontaneously decay (change) into atoms of other elements.

For any given atom, there is a certain probability,  $P$ , that it will undergo radioactive decay in a given amount of time. The half-life,  $\tau$ , is how much time it would take to have a 50% probability of the atom decaying. If you start with  $n$  atoms, after one half-life, half of them ( $0.5n$ ) will have decayed.

If we start with 32 g of  $^{53}\text{Fe}$ , which has a half-life ( $\tau$ ) of 8.5 minutes, we would observe the following:

<b># minutes</b>	0	8.5	17	25.5	34
<b># half lives</b>	0	1	2	3	4
<b>amount left</b>	32 g	16 g	8 g	4 g	2 g

Use this space for summary and/or additional notes:

### Amount of Material Remaining

Most half-life problems in a first-year high school physics course involve a whole number of half-lives and can be solved by making a table like the one above. However, on the AP<sup>®</sup> exam you can expect problems that do not involve a whole number of half-lives, and you need to use the exponential decay equation.

Because  $n$  is decreasing, the number of atoms (and consequently also the mass) remaining after any specific period of time follows the exponential decay function:

$$A = A_0 \left(\frac{1}{2}\right)^n$$

where  $A$  is the amount you have now,  $A_0$  is the amount you started with, and  $n$  is the number of half-lives that have elapsed.

Because the number of half-lives equals the total time elapsed ( $t$ ) divided by the half-life ( $\tau$ ), we can replace  $n = t/\tau$  and rewrite the equation as:

$$A = A_0 \left(\frac{1}{2}\right)^{t/\tau} \quad \text{or} \quad \frac{A}{A_0} = \left(\frac{1}{2}\right)^{t/\tau}$$

If you want to find either  $A$  or  $A_0$ , you can plug the values for  $t$  and  $\tau$  into the above equation.

#### Sample Problem:

Q: If you start with 228 g of <sup>90</sup>Sr, how much would remain after 112.4 years?

A:  $A_0 = 228$  g

$A = A$

$\tau = 28.1$  years (from the "Selected Radioisotopes" table in your reference tables)

$t = 112.4$  years

$$A = A_0 \left(\frac{1}{2}\right)^{t/\tau}$$

$$A = (228) \left(\frac{1}{2}\right)^{112.4/28.1} = (228) \left(\frac{1}{2}\right)^4 = (228) \left(\frac{1}{16}\right) = 14.25 \text{ g}$$

Or, if the decay happens to occur over an integer number of half-lives (as in this example), you can use a chart:

# years	0	28.1	56.2	84.3	112.4
# half lives	0	1	2	3	4
amount left	228 g	114 g	57 g	28.5 g	14.25 g

Use this space for summary and/or additional notes:

## Finding the Time that has Passed

### Integer Number of Half-Lives

If the amount you started with divided by the amount left is an exact power of two, you have an integer number of half-lives and you can just make a table.

### Sample problem:

Q: If you started with 64 g of  $^{131}\text{I}$ , how long would it take until there was only 4 g remaining? The half-life ( $\tau$ ) of  $^{131}\text{I}$  is 8.07 days.

A:  $\frac{64}{4} = 16$  which is a power of 2, so we can simply make a table:

# half lives	0	1	2	3	4
amount remaining	64 g	32 g	16 g	8 g	4 g

From the table, after 4 half-lives, we have 4 g remaining.

The half-life ( $\tau$ ) of  $^{131}\text{I}$  is 8.07 days.

$$8.07 \times 4 = 32.3 \text{ days}$$

Use this space for summary and/or additional notes:

### Non-Integer Number of Half-Lives

If you need to find the elapsed time and it is not an exact half-life, you need to use logarithms.

In mathematics, *the only reason you ever need to use logarithms is when you need to solve for a variable that's in an exponent*. For example, suppose we have the expression of the form  $a^b = c$ .

If  $b$  is a constant, we can solve for either  $a$  or  $c$ , as in the expressions:

$$a^3 = 21 \quad (\sqrt[3]{a^3} = \sqrt[3]{21} = 2.76)$$

$$6^2 = c \quad (6^2 = 36)$$

However, we can't do this if  $a$  and  $c$  are constants and we need to solve for  $b$ , as in the expression:

$$3^b = 17$$

To solve for  $b$ , we need to get  $b$  out of the exponent. We do this by taking the logarithm of both sides:

$$b \log(3) = \log(17)$$

$$b = \frac{\log(17)}{\log(3)} = \frac{1.23}{0.477} = 2.58$$

It doesn't matter which base you use. For example, using  $\ln$  instead of  $\log$  gives the same result:

$$b \ln(3) = \ln(17)$$

$$b = \frac{\ln(17)}{\ln(3)} = \frac{2.83}{1.10} = 2.58$$

We can apply this same logic to the half-life equation:

$$\frac{A}{A_0} = \left(\frac{1}{2}\right)^{t/\tau}$$

$$\log A - \log A_0 = \frac{t}{\tau} \log\left(\frac{1}{2}\right)$$

Use this space for summary and/or additional notes:

**Sample problem:**

Q: If you started with 64 g of  $^{131}\text{I}$ , how long would it take until there was only 5.75 g remaining? The half-life ( $\tau$ ) of  $^{131}\text{I}$  is 8.07 days.

A: We have 5.75 g remaining. However,  $\frac{64}{5.75} = 11.13$ , which is not a power of two.

This means we don't have an integer number of half-lives, so we need to use logarithms:

$$\frac{A}{A_0} = \left(\frac{1}{2}\right)^{\frac{t}{\tau}}$$

$$\log A - \log A_0 = \frac{t}{\tau} \log\left(\frac{1}{2}\right)$$

$$\log 5.75 - \log 64 = \frac{t}{8.07} \log\left(\frac{1}{2}\right)$$

$$0.7597 - 1.8062 = \frac{t}{8.07} (-0.3010)$$

$$-1.0465 = -0.03730 t$$

$$28.1 \text{ days} = t$$

**Homework Problems**

For these problems, you will need to use half-life information from *Table EE*.

*Selected Radioisotopes* on page 624 of your physics reference tables.

1. **(M)** If a lab had 128 g of  $^3\text{H}$  waste 49 years ago, how much of it would be left today? (*Note: you may round off to a whole number of half-lives.*)

Answer: 8 g

Use this space for summary and/or additional notes:

2. **(S)** Suppose you set aside a 20. g sample of  $^{42}\text{K}$  at 5:00pm on a Friday for an experiment, but you are not able to perform the experiment until 9:00am on Monday (64 hours later). How much of the  $^{42}\text{K}$  will be left?

Answer: 0.56 g

3. **(M)** If a school wants to dispose of small amounts of radioactive waste, they can store the materials for ten half-lives, and then dispose of the materials as regular trash.
- a. If we had a sample of  $^{32}\text{P}$ , how long would we need to store it before disposing of it?

Answer: 143 days

- b. If we had started with 64 g of  $^{32}\text{P}$ , how much  $^{32}\text{P}$  would be left after ten half-lives? Approximately what fraction of the original amount would be left?

Answer: 0.063 g; approximately  $\frac{1}{1000}$  of the original amount.

4. **(M)** If the carbon in a sample of human bone contained 30. % of the expected amount of  $^{14}\text{C}$ , approximately how old is the sample?

Answer: 9 950 years

Use this space for summary and/or additional notes: